GRIFOLS

Certificate of Analysis

Product: Cat.No. Lot No.: Expiration date:

Extended IV Control 213286 QCX022007 2022-05-14

Origin: human erythrocytes from blood group A2B, A, B and O

Manufacturing Date: 28 MAR 2022 Preservative: Neomycin & Chloramphenicol

Storage: 2-8 °C

Manufacturing size: 4 x 6 ml

Blood group testing by DG Gel	Blood group A ₂ B CcD.Ee, K positive	Blood group A ₂ B CcD.Ee, K positive
Blood group testing by MDmulticard (Lateral Flow)	Blood group AB CcD.Ee, K positive	Blood group AB CcD.Ee, K positive
Blood group testing by tube technique	Blood group AB CcD.Ee, K positive	Blood group AB CcD.Ee, K positive
Reverse-Typing by DG Gel	A ₁ cells: negative A ₂ cells: negative B cells: negative O cells: negative	A ₁ cells: negative A ₂ cells: negative B cells: negative O cells: negative
Reverse-Typing by tube technique	A ₁ cells: negative A ₂ cells: negative B cells: negative O cells: negative	A ₁ cells: negative A ₂ cells: negative B cells: negative O cells: negative
Direct Antiglobulin Test by DG Gel at release	negative	negative
Direct Antiglobulin Test by tube technique at release	negative	negative
IAT Test by DG Gel with Screen-Cyte cells	Screen-Cyte cells: negative	Screen-Cyte cells: negative
IAT test by tube technique with Screen-Cyte cells	Screen-Cyte cells: negative	Screen-Cyte cells: negative
Determination of Hematocrit	15.0 ± 2.0 %	13.4 %
Photometry at 540nm	≤ 30.0 mg/dl of Hemoglobin	3.65 mg/dl of Hemoglobin
	Blood group testing by MDmulticard (Lateral Flow) Blood group testing by tube technique Reverse-Typing by DG Gel Reverse-Typing by tube technique Direct Antiglobulin Test by DG Gel at release Direct Antiglobulin Test by tube technique at release IAT Test by DG Gel with Screen-Cyte cells IAT test by tube technique with Screen-Cyte cells Determination of Hematocrit	Blood group testing by MDmulticard (Lateral Flow) Blood group testing by tube technique Reverse-Typing by DG Gel Reverse-Typing by tube R

Tube QCX2, lot QCX222007			
Specificity	Blood group testing by DG Gel	Blood group A CC ^w D.ee, Fy ^a negative	Blood group A CC ^w D.ee, Fy ^a negative
	Blood group testing by MDmulticard (Lateral Flow)	Blood group A CC ^w D.ee	Blood group A CC ^w D.ee
	Blood group testing by tube technique	Blood group A CC ^w D.ee, Fy ^a negative	Blood group A CC ^w D.ee, Fy ^a negative
	Reverse-Typing by DG Gel	A ₁ cells: negative A ₂ cells: negative B cells: positive O cells: negative	A ₁ cells: negative A ₂ cells: negative B cells: positive O cells: negative
	Reverse-Typing by tube technique	A ₁ cells: negative A ₂ cells: negative B cells: positive O cells: negative	A ₁ cells: negative A ₂ cells: negative B cells: positive O cells: negative
	Direct Antiglobulin Test by DG Gel at release	negative	negative
	Direct Antiglobulin Test by tube technique at release	negative	negative
Fy ^a Antibody screening	IAT test by DG Gel with Screen-Cyte	Screening cells Fy ^a positive: positive Screening cells Fy ^a negative: negative	Screening cells Fy ^a positive: positive Screening cells Fy ^a negative: negative
	IAT test by tube technique with Screen-Cyte cells	Screening cells Fy ^a positive: positive Screening cells Fy ^a negative: negative	Screening cells Fy ^a positive: positive Screening cells Fy ^a negative: negative
Concentration of suspension	Determination of Hematocrit		13.4 %
Hemolysis after filling	Photometry at 540nm	≤ 30 mg/dl of Hemoglobin	3.65 mg/dl of Hemoglobin
Tb. 00V2 Int 00V222007			
Tube QCX3, lot QCX322007 Specificity	Blood group testing by DG Gel	Blood group B ccD.EE	Blood group B ccD.EE
	Blood group testing by MDmulticard (Lateral Flow)	Blood group B ccD.EE	Blood group B ccD.EE
	Blood group testing by tube technique	Blood group B ccD.EE	Blood group B ccD.EE
	Reverse-Typing by DG Gel	A ₁ cells: positive A ₂ cells: positive B cells: negative O cells: negative	A ₁ cells: positive A ₂ cells: positive B cells: negative O cells: negative
	Reverse-Typing by tube technique	A ₁ cells: positive A ₂ cells: positive B cells: negative O cells: negative	A ₁ cells: positive A ₂ cells: positive B cells: negative O cells: negative
	Direct Antiglobulin Test by DG Gel at release	negative	negative
	Direct Antiglobulin Test by tube technique at release	negative	negative
Antibody screening	IAT Test by DG Gel with Screen-Cyte cells	Screen-Cyte cells: negative	Screen-Cyte cells: negative
	IAT test by tube technique with Screen-Cyte cells	Screen-Cyte cells: negative	Screen-Cyte cells: negative
Concentration of suspension	Determination of Hematocrit	15.0 ± 2.0 %	13.3 %
Hemolysis after filling	Photometry at 540nm	≤ 30 mg/dl of Hemoglobin	3.65 mg/dl of Hemoglobin

Tube QCX4, lot QCX422007			
Specificity	Blood group testing by DG Gel	Blood group O ccddee, K negative	Blood group O ccddee, K negative
	Blood group testing by MDmulticard (Lateral Flow)	Blood group O ccddee, K negative	Blood group O ccddee, K negative
	Blood group testing by tube technique	Blood group O ccddee, K negative	Blood group O ccddee, K negative
	Reverse-Typing by DG Gel	A ₁ cells: positive A ₂ cells: positive B cells: positive O cells: negative	A ₁ cells: positive A ₂ cells: positive B cells: positive O cells: negative
	Reverse-Typing by tube technique	A ₁ cells: positive A ₂ cells: positive B cells: positive O cells: negative	A ₁ cells: positive A ₂ cells: positive B cells: positive O cells: negative
	Direct Antiglobulin Test by DG Gel at release	negative	negative
	Direct Antiglobulin Test by tube technique at release	negative	negative
D Antibody screening	IAT test by DG Gel IgG	Ror cells: minimum weak positive R ₁ R ₁ cells: minimum weak positive R ₂ R ₂ cells: minimum weak positive rr cells: negative	Ror cells: minimum weak positive R ₁ R ₁ cells: minimum weak positive R ₂ R ₂ cells: minimum weak positive rr cells: negative
	IAT test by tube technique	Ror cells: minimum weak positive R ₁ R ₁ cells: minimum weak positive R ₂ R ₂ cells: minimum weak positive rr cells: negative	Ror cells: minimum weak positive R ₁ R ₁ cells: minimum weak positive R ₂ R ₂ cells: minimum weak positive rr cells: negative
Concentration of suspension	Determination of Hematocrit		13.4 %
Hemolysis after filling	Photometry at 540nm	≤ 30 mg/dl of Hemoglobin	2.92 mg/dl of Hemoglobin

The product passed the internal Quality Control procedure.

All human source material used in the manufacture of this product were tested and found negative for HBsAg and antibodies to HCV, HIV-1/ HIV-2.

Date: 31 March 2022

Sascha Feuerhahn Technical Director

Medion Grifols Diagnostics AG in Düdingen, Switzerland, is certified according to EN ISO 13485.